



Certificate of Analysis

www.tocris.com

Product Name: [D-Ala²]-Deltorphin II Catalog No.: 1180 Batch No.: 25

CAS Number: 122752-16-3

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: $C_{38}H_{54}N_8O_{10}$

Batch Molecular Weight: 782.89

Physical Appearance: White lyophilised solid

Counter Ion: TFA

Solubility: Soluble to 1 mg/ml in water

Storage: Store at -20°C

Peptide Sequence: Tyr-D-Ala-Phe-Glu-Val-Val-Gly-NH₂

2. ANALYTICAL DATA

HPLC: Shows 95.3% purity

Mass Spectrum: Consistent with structure

3. AMINO ACID ANALYSIS DATA

Ala	1.00	0.98	Lys		
Arg			Met		
Asx			Phe	1.00	1.03
Cys			Pro		
Glx	1.00	1.01	Ser		
Gly	1.00	1.02	Thr		
His			Trp		
lle			Tyr	1.00	0.96
Leu			Val	2.00	1.80

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use



Product Name:

Product Information

Print Date: Apr 18th 2024

25

www.tocris.com

CAS Number: 122752-16-3

Description:

[D-Ala²]-Deltorphin II is a selective peptide agonist for the δ opioid receptor. Antinociceptive in vivo.

[D-Ala²]-Deltorphin II

Physical and Chemical Properties:

Batch Molecular Formula: $C_{38}H_{54}N_8O_{10}$ Batch Molecular Weight: 782.89

Physical Appearance: White lyophilised solid

Peptide Sequence:

Tyr-D-Ala-Phe-Glu-Val-Val-Gly-NH2

Storage: Store at -20°C

Solubility & Usage Info:

Soluble to 1 mg/ml in water

This product is supplied in lyophilized form. It may appear as a solid, gel or film and be very hard to visualize. Solutions should be made by adding solvent directly to the vial. The vial should then be vortexed vigorously to ensure the product has completely dissolved.

Catalog No.: 1180

Counter Ion: TFA

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Peptides in solution are much less stable than in lyophilized form. This is especially true for peptides whose sequences contain amino acids such Cys, Met,Trp, Asn, Gln, and N-terminal Glu.

Therefore we recommend storing peptides in solution for as short a time as possible. Avoid repeated freeze thaw cycles by dividing the peptide solution into aliquots and storing the aliquots at -20°C. Any portion of an aliquot unused after thawing should be discarded.

Peptides stored in solution can occasionally be susceptible to bacterial degradation. We recommend using sterile solutions or passing the peptide solution through a 0.2 μ m filter to remove potential bacterial contamination whenever possible.

References:

Sanchez-Blazquez (1998) Delta opioid receptor subtypes activate inositol-signalling pathways in the production of antinociception. J.Pharmacol.Exp.Ther. **285** 820. PMID: 9580632.

Thomas et al (1997) Structure-activity relationships of a series of [D-Ala²] deltorphin I and II analogues; in vitro blood-brain barrier permeability and stability. J.Pharmacol.Exp.Ther. **281** 817. PMID: 9152390.

Mattia et al (1991) Lack of antinociceptive cross-tolerance between [D-Pen², D-Pen⁵]enkephalin and [D-Ala²]deltorphin II in mice: evidence for delta receptor subtypes. J.Pharmacol.Exp.Ther. **258** 583. PMID: 1650835.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use