

Product Name: Pepstatin A

Catalog No.: 1190


Batch No.: 22

CAS Number: 26305-03-3

EC Number: 247-600-0

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: C₃₄H₆₃N₅O₉
Batch Molecular Weight: 685.91
Physical Appearance: White solid
Solubility: Soluble to 1 mg/ml in ethanol with gentle warming
Storage: Desiccate at -20°C
Peptide Sequence:



2. ANALYTICAL DATA

HPLC: Shows 99.1% purity

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

Product Name: Pepstatin A

Catalog No.: 1190

Batch No.: 22

CAS Number: 26305-03-3

EC Number: 247-600-0

Description:

Irreversible inhibitor of aspartic proteases. Inhibits lysosomal proteases and interferes with autolysosomal digestion when used in combination with E 64d (Cat. No. 4545).


Physical and Chemical Properties:

Batch Molecular Formula: C₃₄H₆₃N₅O₉

Batch Molecular Weight: 685.91

Physical Appearance: White solid

Peptide Sequence:



Storage: Desiccate at -20°C

Solubility & Usage Info:

Soluble to 1 mg/ml in ethanol with gentle warming

This product is supplied as a lyophilized solid and may be very hard to visualize. Solutions should be made by adding solvent directly to the vial. The vial should then be vortexed vigorously to ensure the product has completely dissolved.

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Peptides in solution are much less stable than in lyophilized form. This is especially true for peptides whose sequences contain amino acids such as Cys, Met, Trp, Asn, Gln, and N-terminal Glu.

Therefore we recommend storing peptides in solution for as short a time as possible. Avoid repeated freeze thaw cycles by dividing the peptide solution into aliquots and storing the aliquots at -20°C. Any portion of an aliquot unused after thawing should be discarded.

Peptides stored in solution can occasionally be susceptible to bacterial degradation. We recommend using sterile solutions or passing the peptide solution through a 0.2 µm filter to remove potential bacterial contamination whenever possible.

References:

Sato et al (2007) Autophagy is activated in colorectal cancer cells and contributes to the tolerance to nutrient deprivation. *Cancer Res.* **67** 9677. PMID: 17942897.

Marciniszyn et al (1976) Mode of inhibition of acid proteases by pepstatin. *J.Biol.Chem.* **251** 7088. PMID: 993206.

Merck Index **12** 7290.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com

info@bio-techne.com

techsupport@bio-techne.com

North America

Tel: (800) 343 7475

China

info.cn@bio-techne.com

Tel: +86 (21) 52380373

Europe Middle East Africa

Tel: +44 (0)1235 529449

Rest of World

www.tocris.com/distributors

Tel:+1 612 379 2956