

Product Name: Lovastatin

Catalog No.: 1530


Batch No.: 1

CAS Number: 75330-75-5

IUPAC Name: (2S)-(1S,3R,7S,8S,8aR)-1,2,3,7,8,8a-Hexahydro-3,7-dimethyl-8-[2-[(2R,4R)-tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl]ethyl]-1-naphthalenyl-2-methyl butanoate

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: C₂₄H₃₆O₅
Batch Molecular Weight: 404.54
Physical Appearance: White solid
Solubility: ethanol to 50 mM with gentle warming
DMSO to 100 mM
Storage: Store at RT
Batch Molecular Structure:



2. ANALYTICAL DATA

TLC: R_f = 0.48 (Dichloromethane:Methanol [10:1])
Melting Point: At 178°C
HPLC: Shows 98.5% purity
¹H NMR: Consistent with structure

Microanalysis:

	Carbon	Hydrogen	Nitrogen
Theoretical	71.26	8.97	
Found	71.3	9.01	

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

Product Name: Lovastatin

Catalog No.: 1530

Batch No.: 1

CAS Number: 75330-75-5

IUPAC Name: (2S)-(1S,3R,7S,8S,8aR)-1,2,3,7,8,8a-Hexahydro-3,7-dimethyl-8-[2-[(2R,4R)-tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl]ethyl]-1-naphthalenyl-2-methyl butanoate

Description:

Potent, competitive inhibitor of HMG-CoA reductase ($K_i = 0.6$ nM) therefore decreases cholesterol biosynthesis, in vitro and in vivo. Indirectly inhibits Rho-GTPase activity by depleting intracellular pool of isoprene residues. Decreases CDK2, 4, 6 and cyclin E levels and induces G1 arrest and apoptosis in tumor cell lines in vitro. Inactive lactam prodrug of lovastatin hydroxy acid, naturally bioactivated in vivo following oral administration.

Physical and Chemical Properties:


Batch Molecular Formula: C₂₄H₃₆O₅

Batch Molecular Weight: 404.54

Physical Appearance: White solid

Minimum Purity: >98%

Batch Molecular Structure:



Storage: Store at RT

Solubility & Usage Info:

ethanol to 50 mM with gentle warming

DMSO to 100 mM

PLEASE NOTE - Lovastatin is an inactive lactam prodrug of lovastatin hydroxy acid the active form of the compound. Lovastatin administered orally is naturally bioactivated by the liver and requires no further modification. If lovastatin is to be administered by another route or used in vitro then it requires manual activation by treatment with NaOH. To activate the compound dissolve 50mg in 1ml of warm (50°C) ethanol and add 0.813ml of 1N NaOH. Leave for 30 minutes to allow conversion of the lovastatin to the active acid form, the compound may be stored at -20°C in this format for up to 1 month. Adjust pH to 7.2 with small quantities of 1N HCl prior to use.

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Ziegler et al (2017) Rho inhibition by lovastatin affects apoptosis and DSB repair of primary human lung cells *in vitro* and lung tissue *in vivo* following fractionated irradiation. *Cell Death Dis.* **8** e2978. PMID: 28796249 .

Park et al (1999) Lovastatin-induced inhibition of HL-60 cell proliferation via cell cycle arrest and apoptosis. *Anticancer Res.* **19** 3133. PMID: 10652602.

Alberts (1988) Discovery, biochemistry and biology of lovastatin. *Am.J.Cardiol.* **62** 10J. PMID: 3055919.

Alberts et al (1980) Mevinolin: a highly potent competitive inhibitor of hydroxymethylglutaryl-coenzyme A reductase and a cholesterol-lowering agent. *Proc.Natl.Acad.Sci.U.S.A.* **77** 3957. PMID: 6933445.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com

info@bio-techne.com

techsupport@bio-techne.com

North America

Tel: (800) 343 7475

China

info.cn@bio-techne.com

Tel: +86 (21) 52380373

Europe Middle East Africa

Tel: +44 (0)1235 529449

Rest of World

www.tocris.com/distributors

Tel:+1 612 379 2956