TOCRIS a biotechne brand

Certificate of Analysis

www.tocris.com

Catalog No.: 3905

Print Date: Oct 8th 2019


Batch No.: 2

Product Name: NAADP tetrasodium salt

IUPAC Name: Nicotinic acid adenine dinucleotide phosphate tetrasodium salt

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: Batch Molecular Weight: Physical Appearance: Solubility: Storage: Batch Molecular Structure: C₂₁H₂₃N₆Na₄O₁₈P₃ 832.32 White solid water to 50 mM Store at -20°C



2. ANALYTICAL DATA

HPLC: Mass Spectrum:

Shows 99.0% purity Consistent with structure

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

TOCRIS a biotechne brand

Product Information

Batch No.: 2

www.tocris.com

Product Name: NAADP tetrasodium salt

IUPAC Name:

Nicotinic acid adenine dinucleotide phosphate tetrasodium salt

Description:


Ca²⁺ mobilizing agent. Initiates Ca²⁺ release via type 1 ryanodine receptor (RyR1) activation. Also regulates Ca²⁺ release from intracellular stores distinct from the endoplasmic reticulum (ER); displays affinity for two-pore channels (TPCs) which release Ca²⁺ from acidic organelles.

Physical and Chemical Properties:

Batch Molecular Formula: $C_{21}H_{23}N_6Na_4O_{18}P_3$ Batch Molecular Weight: 832.32 Physical Appearance: White solid

Minimum Purity: >98%

Batch Molecular Structure:



Storage: Store at -20°C

Solubility & Usage Info:

water to 50 mM

Solutions of NAADP should ideally be prepared and used as quickly as possible. Solutions of 10 mg/mLstored at 2-8°C at neutral pH are expected to lose 2-5% activity per week. Solutions stored at -20°C are expected to lose between 2-5% activity after six months. Lyophilized NAADP shows a loss of activity of less than 2% after 12 months when stored at -20°C

Catalog No.: 3905

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Calcraft *et al* (2009) NAADP mobilizes calcium from acidic organelles through two-pore channels. Nature **459** 596. PMID: 19387438. **Dammerman** *et al* (2009) NAADP-mediated Ca²⁺ signaling via type 1 ryanodine receptor in T cells revealed by a synthetic NAADP agonist. Proc.Natl.Acad.Sci. **106** 10678.

Galion (2006) NAADP, a new intracellular messenger that mobilizes Ca2+ from acidic stores. Biochem.Soc.Transactions 34 922.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com	North America	China	Europe Middle East Africa	Rest of World
info@bio-techne.com techsupport@bio-techne.com	Tel: (800) 343 7475	info.cn@bio-techne.com Tel: +86 (21) 52380373	Tel: +44 (0)1235 529449	www.tocris.com/distributors Tel:+1 612 379 2956