

Certificate of Analysis

Print Date: Sep 6th 2024

www.tocris.com

Product Name: LDV FITC Catalog No.: 4577 Batch No.: 8

CAS Number: 1207610-07-8

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: C₆₉H₈₁N₁₁O₁₇S

Batch Molecular Weight: 1368.54

Physical Appearance: Yellow lyophilised solid

Counter Ion: TFA

Solubility: Soluble to 1 mg/ml in PBS (pH 7.4)

Storage: Store at -20°C

Peptide Sequence:

2. ANALYTICAL DATA

HPLC: Shows 94.3% purity

Mass Spectrum: Consistent with structure

3. AMINO ACID ANALYSIS DATA

Amino Acid Theoretical Actual Amino Acid Theoretical Actual

Ala	2.00	1.97	Lys	1.00	Not Detected
Arg			Met		
Asx	1.00	1.02	Phe		
Cys			Pro	1.00	0.98
Glx			Ser		
Gly			Thr		
His			Trp		
lle			Tyr		
Leu	1.00	1.01	Val	1.00	1.01

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use



Product Information

Print Date: Sep 6th 2024

www.tocris.com

Product Name: LDV FITC Catalog No.: 4577 8

CAS Number: 1207610-07-8

Description:

LDV FITC is a fluorescent ligand that binds to the $\alpha_4\beta_1$ integrin (VLA-4) with high affinity (K_d values are 0.3 nM and 12 nM for binding to U937 cells in the presence and absence of Mn²+ respectively). Used to detect VLA-4 affinity and conformational changes. Wavelengths are compatible with use as an acceptor dye in TR-FRET assays, together with the CoraFluor 1 (Cat. No. 7920) TR-FRET donor.

Physical and Chemical Properties:

Batch Molecular Formula: C₆₉H₈₁N₁₁O₁₇S Batch Molecular Weight: 1368.54

Physical Appearance: Yellow lyophilised solid

Peptide Sequence:

Storage: Store at -20°C

Solubility & Usage Info:

Soluble to 1 mg/ml in PBS (pH 7.4)

This product is supplied in lyophilized form. It may appear as a solid, gel or film and be very hard to visualize. Solutions should be made by adding solvent directly to the vial. The vial should then be vortexed vigorously to ensure the product has completely dissolved.

Counter Ion: TFA

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Peptides in solution are much less stable than in lyophilized form. This is especially true for peptides whose sequences contain amino acids such Cys, Met,Trp, Asn, Gln, and N-terminal Glu.

Therefore we recommend storing peptides in solution for as short a time as possible. Avoid repeated freeze thaw cycles by dividing the peptide solution into aliquots and storing the aliquots at -20°C. Any portion of an aliquot unused after thawing should be discarded.

Peptides stored in solution can occasionally be susceptible to bacterial degradation. We recommend using sterile solutions or passing the peptide solution through a 0.2 μ m filter to remove potential bacterial contamination whenever possible.

References:

Njus *et al* (2009) Conformational mAb as a tool for integrin ligand discovery. Assay Drug Dev.Technol. **7** 507. PMID: 19754304. **Chigaev** *et al* (2003) FRET detection of cellular α₄-integrin conformational activation. Biophys.J. **85** 3951. PMID: 14645084.

Chigaev *et al* (2001) Real time analysis of the affinity regulation of alpha 4-integrin. The physiologically activated receptor is intermediate in affinity between resting and Mn²⁺ or antibody activation. J.Biol.Chem. **276** 48670. PMID: 11641394.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use